



**DOMESTIC & COMMERCIAL  
PEST CONTROL**

Same Family Business Since 1964

**fumapest.com.au**

**1300 241 500**

**Pest Control of Birds • Pigeons • Starlings • Indian Mynah Birds**



• **Some birds are real pests** Pigeons, starlings and indian mynah birds have adapted to live in or around buildings occupied by humans - they survive largely by scavenging for food, water, materials and shelter provided by human habitat.

• **Pigeons** often roost on building ledges, causing

defacement and health risks from their droppings.

• **Starlings - Indian Mynah - Sparrows** often cause big problems to home owners when they nest inside roof voids, causing bird mites to enter the premises. The bites of bird mites cause severe skin irritation and blotching.

**Pigeon Control**

• **Deterrent systems - pigeons** - various long term bird control deterrent products can be used where pigeons are a problem - installation of plastic or stainless steel spikes on roosting ledges - post and spring-wire barriers - bird netting to exclude pigeons from particular areas.

• **Shooting** of pigeons may be allowed if approved by the National Parks & Wildlife Service and the local police authorities. Live trapping and humane disposal of pigeons, starlings, indian mynah birds and other bird pests is also permitted in certain circumstances.

**Bird Mite Infestation & Control**

• **Bird mite infestations** are common throughout NSW during the warmer summer months - the bird mite (sometimes wrongly called "bird lice") most often come from **starlings, indian mynah or sparrows** nesting inside roof voids or other areas of a building.

• **Severe skin reaction** - the bites of bird mites are 10 times worse than fleas and cause severe skin irritation and blotching, particularly to children or people with sensitive skin.

• **Pest control for bird mites** - firstly, FUMAPEST will conduct a thorough inspection of the building and surrounds - a quotation supplied for immediate eradication of the bird mites - and a quotation for closing off possible entry points for the birds into the roof void, if applicable.

• **Safer insecticides** FUMAPEST uses the safer, but more expensive, synthetic pyrethroid insecticide to immediately treat the bird nest

containing the bulk of the bird mites, prior to removal of the bird nesting materials. The surrounding roof void areas is then treated with a permethrin dust.

• **Inside the home** - any bird mite infested areas in the living quarters of the home is treated with natural pyrethrins, as applied through a fogging machine that disperses the natural pyrethrin insecticide to penetrate into nooks and crannies where the bird mite may congregate. Essential for immediate relief.

• **To avoid on-going problems** it is also essential to block off all possible entry points to ensure the birds cannot return. If entry points are left unchanged, then the Starlings or Indian Mynah birds will most likely return to the same location to build another nest to raise their young. FUMAPEST uses a range of bird proofing materials designed specifically for bird proofing of domestic and commercial premises.

**RING FUMAPEST ON 1300 241 500 FOR PEST CONTROL OF BIRDS**

Termites - Termite Control - Domestic Pest Control - Commercial Pest Control - Consumer Advice

FUMAPEST Termite & Pest Control © copyright 1995 - 2020